

# Cycle Spectra of Hamiltonian Graphs

Kevin G. Milans<sup>†</sup>, Florian Pfender<sup>‡</sup>, Dieter Rautenbach<sup>§</sup>,  
Friedrich Regen<sup>§</sup>, and Douglas B. West<sup>¶</sup>

March 23, 2012

## Abstract

We prove that every Hamiltonian graph with  $n$  vertices and  $m$  edges has cycles with more than  $\sqrt{p} - \frac{1}{2} \ln p - 1$  different lengths, where  $p = m - n$ . For general  $m$  and  $n$ , there exist such graphs having at most  $2 \lceil \sqrt{p+1} \rceil$  different cycle lengths.

**Keywords:** cycle, cycle spectrum, Hamiltonian graph, Hamiltonian cycle.

## 1 Introduction

The *cycle spectrum* of a graph  $G$  is the set of lengths of cycles in  $G$ . A cycle containing all vertices of a graph is a *spanning* or *Hamiltonian cycle*, and a graph having such a cycle is a *Hamiltonian graph*. An  $n$ -vertex graph is *pancyclic* if its cycle spectrum is  $\{3, \dots, n\}$ . Our graphs have no loops or multiple edges. A graph is  $k$ -regular if every vertex has degree  $k$  (that is,  $k$  incident edges).

Interest in cycle spectra arose from Bondy's "Metaconjecture" (based on [3]) that sufficient conditions for the existence of Hamiltonian cycles usually also imply that a graph is pancyclic, with possibly a small family of exceptions. For example, Bondy [3] showed that the sufficient condition on  $n$ -vertex graphs due to Ore [16] (the degrees of any two nonadjacent vertices sum to at least  $n$ ) implies also that  $G$  is pancyclic or is the complete bipartite

---

<sup>†</sup>Mathematics Dept., University of South Carolina, milans@math.sc.edu.

<sup>‡</sup>Institut für Mathematik, Universität Rostock, Rostock, Germany, florian.pfender@uni-rostock.de.

<sup>§</sup>Institut für Optimierung und Operations Research, Universität Ulm, Ulm, Germany, dieter.rautenbach@uni-ulm.de and friedrich.regen@uni-ulm.de; supported in part by the Deutsche Forschungsgemeinschaft DFG within the project "Cycle Spectra of Graphs".

<sup>¶</sup>Department of Mathematics, University of Illinois, Urbana, IL 61801, USA, west@math.uiuc.edu; supported in part by the National Security Agency under Awards H98230-06-1-0065 and H98230-10-1-0065.

graph  $K_{\frac{n}{2}, \frac{n}{2}}$ . Schmeichel and Hakimi [13] showed that if a spanning cycle in an  $n$ -vertex graph  $G$  has consecutive vertices with degree-sum at least  $n$ , then  $G$  is pancyclic or bipartite or omits only  $n - 1$  from the cycle spectrum, the latter occurring only when the degree-sum is exactly  $n$ . Bauer and Schmeichel [1] used this to give unified proofs that the conditions for Hamiltonian cycles due to Bondy [4], Chvátal [5], and Fan [9] also imply that a graph is pancyclic, with small families of exceptions. Further results about the cycle spectrum under degree conditions on selected vertices in a spanning cycle appear in [10] and [14].

At the 1999 conference “Paul Erdős and His Mathematics”, Jacobson and Lehel proposed the opposite question: *When sufficient conditions for spanning cycles are relaxed, how small can the cycle spectrum be if the graph is required to be Hamiltonian?* For example, consider regular graphs. Bondy’s result [3] implies that  $\lceil n/2 \rceil$ -regular graphs other than  $K_{\frac{n}{2}, \frac{n}{2}}$  are pancyclic. On the other hand, 2-regular Hamiltonian graphs have only one cycle length. For  $3 \leq k \leq \lceil n/2 \rceil - 1$ , Jacobson and Lehel asked for the minimum size of the cycle spectrum of a  $k$ -regular  $n$ -vertex Hamiltonian graph, particularly when  $k = 3$ .

Let  $s(G)$  denote the size of the cycle spectrum of a graph  $G$ . At the SIAM Meeting on Discrete Mathematics in 2002, Jacobson announced that he, Gould, and Pfender had proved  $s(G) \geq c_k n^{1/2}$  for  $k$ -regular graphs with  $n$  vertices. Others later independently obtained similar bounds, without seeking to optimize  $c_k$ . For an upper bound, Jacobson and Lehel constructed the 3-regular example below with only  $n/6 + 3$  distinct cycle lengths (when  $n \equiv 0 \pmod{6}$  and  $n > 6$ ), and they generalized it to the upper bound  $\frac{n}{2} \frac{k-2}{k} + k$  for  $k$ -regular graphs.

**Example 1** When  $k = 3$  and 6 divides  $n$  (with  $n > 6$ ), take  $n/6$  disjoint copies of  $K_{3,3}$  in a cyclic order, with vertex sets  $V_1, \dots, V_{n/6}$ . Remove one edge from each copy and replace it by an edge to the next copy to restore 3-regularity. A cycle of length different from 4 or 6 must visit each  $V_i$ , and in each  $V_i$  it uses 4 or 6 vertices. Hence the cycle lengths are 4, 6, and each even integer from  $2n/3$  through  $n$ . For the generalization, use  $K_{k,k}$  instead of  $K_{3,3}$ .  $\square$

A related problem is the conjecture of Erdős [7] that  $s(G) \geq \Omega(d^{\lfloor (g-1)/2 \rfloor})$  when  $G$  has girth  $g$  and average degree  $d$ . Erdős, Faudree, Rousseau, and Schelp [8] proved the conjecture for  $g = 5$ . Sudakov and Verstraëte [15] proved the full conjecture in a stronger form, obtaining  $\frac{1}{8} (d^{\lfloor (g-1)/2 \rfloor})$  consecutive even integers in the cycle spectrum for graphs with fixed girth  $g$  and average degree  $48(d + 1)$ . Gould, Haxell, and Scott [11] proved a similar result: for  $c > 0$ , there is a constant  $k_c$  such that for sufficiently large  $n$ , the cycle spectrum of every  $n$ -vertex graph  $G$  having minimum degree at least  $cn$  and longest even cycle length  $2l$  contains all even integers from 4 up to  $2l - k_c$  (see also [2]).

Prior arguments for lower bounds on  $s(G)$  when  $G$  is regular and Hamiltonian used only the number of edges, not regularity. Suppose that  $G$  has  $n$  vertices and  $m$  edges. The

coefficient  $c$  in a general lower bound of the form  $s(G) \geq \sqrt{c(m-n)}$  cannot exceed 1, since  $s(K_{\frac{n}{2}, \frac{n}{2}}) = \sqrt{m-n+1}$ . We give a construction for  $m \leq n^2/4$  that is far from regular.

**Example 2** For  $t \leq n/2$ , form a graph  $G$  by replacing one edge of  $K_{t,t}$  with a path having  $n-2t$  internal vertices;  $G$  has  $n$  vertices and  $m$  edges, where  $m = t^2 - 2t + n \leq n^2/4$ . The cycle spectrum of  $G$  consists of the  $t-1$  even numbers in  $\{4, \dots, 2t\}$  and the  $t-1$  numbers from  $n-2t+4$  to  $n$  having the same parity as  $n$ . Thus  $s(G) \leq 2(t-1) = 2\sqrt{m-n+1}$ . Equality holds when  $t \leq \lceil n/4 \rceil$ , but when  $\lceil n/4 \rceil < t \leq n/2$  and  $n$  is even the two sets of  $t-1$  numbers overlap. They overlap more as  $m$  increases, becoming the same set when  $m = n^2/4$ , and indeed  $s(K_{\frac{n}{2}, \frac{n}{2}}) = \sqrt{m-n+1}$ .

Deleting edges cannot enlarge the cycle spectrum. Hence in general we can let  $t = \lceil \sqrt{m-n+1} \rceil + 1$ , apply the construction above for  $n$  and  $t$ , and discard edges to wind up with  $m$  edges and  $s(G) \leq 2 \lceil \sqrt{m-n+1} \rceil$ .  $\square$

Bondy [3] showed that every Hamiltonian graph with more than  $n^2/4$  edges is pancyclic. Thus the lower bound on  $s(G)$  jumps to  $n-2$  when  $m$  exceeds  $n^2/4$ . At  $m = n^2/4$ , the size of the spectrum of  $K_{n/2, n/2}$  is only  $n/2 - 1$ . For  $n$ -vertex Hamiltonian bipartite graphs (with  $n > 6$ ), Entringer and Schmeichel [6] proved that  $m > n^2/8$  suffices to make the graph *bipancyclic*, meaning that it has cycles of all  $n/2 - 1$  even lengths.

In the construction of Example 2, the two segments overlap to yield bipancyclic graphs when  $m$  exceeds  $n^2/16 + n/2$ . The result of [6] implies that the construction is optimal among Hamiltonian bipartite graphs when  $m$  exceeds  $n^2/8$ , but whether this also holds for Hamiltonian non-bipartite graphs is unknown. It is also unknown whether there are non-bipancyclic constructions (bipartite or not) when  $n^2/16 + n/2 < m \leq n^2/8$ .

When  $m < n^2/4$ , the construction of Example 2 remains a candidate for a graph having the smallest cycle spectrum among Hamiltonian graphs with  $n$  vertices and  $m$  edges. We do know of one exception: when  $(n, m) = (14, 21)$ , the cycle spectrum of the Heawood graph (incidence graph of the projective plane of order 2) is smaller.

Our main result for the cycle spectra of  $n$ -vertex Hamiltonian graphs with  $m$  edges is that  $s(G) > \sqrt{p} - \frac{1}{2} \ln p - 1$ , where  $p = m - n$ .

## 2 The Lower Bound

A path with endpoints  $x$  and  $y$  is an  $x, y$ -*path*. A *chord* of a path (or cycle)  $P$  in a graph is an edge of the graph not in  $P$  whose endpoints are in  $P$ , and the *length* of the chord is the distance in  $P$  between its endpoints. In a path with vertices  $v_1, \dots, v_n$  in order, two chords  $v_a v_c$  and  $v_b v_d$  *overlap* if  $a < b < c < d$ .

**Lemma 3** *If a graph  $G$  consists of an  $x, y$ -path  $P$  and  $h$  pairwise-overlapping chords of length  $l$ , then  $G$  contains  $x, y$ -paths having at least  $h - 1$  distinct lengths. Having only  $h - 1$  lengths requires  $l$  odd,  $h \geq (l + 3)/2$ , and chords starting at  $h$  consecutive vertices along  $P$ .*

**Proof.** The claim is trivial for  $h = 1$ ; assume  $h \geq 2$ . Let  $n$  be the length of  $P$ . Let  $e_1, \dots, e_h$  be the chords in order of appearance along  $P$  from  $x$  to  $y$ . Let  $d_i$  be the distance along  $P$  from the first endpoint of  $e_{i-1}$  to the first endpoint of  $e_i$ , for  $2 \leq i \leq h$ .

Let  $P_{i,j}$  be the unique  $x, y$ -path using exactly two chords  $e_i$  and  $e_j$ , along with edges of  $P$ . Let  $p_j$  be the length of  $P_{1,j}$ , for  $2 \leq j \leq h$ . Note that  $p_j = p_{j-1} - 2d_j$  for  $3 \leq j \leq h$ . The  $h - 1$  paths  $P_{1,2}, \dots, P_{1,h}$  have distinct lengths, which proves the first statement.

The length of  $P_{1,2}$  is  $n - 2d_2 + 2$ . Thus the full path  $P$  provides an additional length unless  $d_2 = 1$ . If  $d_j > 1$  for any larger  $j$ , then the length of  $P_{2,j}$  is strictly between  $p_{j-1}$  and  $p_j$ . Hence an extra length arises unless the chords start at consecutive vertices along  $P$ .

In the remaining case, the  $h - 1$  lengths we have found are  $n, n - 2, \dots, n - 2h + 4$ . The length of any  $x, y$ -path that uses exactly one chord is  $n - l + 1$ . To avoid generating a new length, it must be that  $l$  is odd and  $2h - 4 \geq l - 1$ .  $\square$

**Definition 4** Let  $G$  be a graph consisting of an  $n$ -cycle  $C$  plus  $q$  chords of length  $l$ , where  $l < n/2$ . Specify a forward direction along  $C$ . Let  $C[u, v]$  denote the subpath of  $C$  traversed by moving forward from  $u$  to  $v$  along  $C$ . When  $uv$  is a chord of length  $l$  and  $C[u, v]$  has length  $l$ , we say that  $u$  is its *start*,  $v$  is its *end*, and  $uv$  *covers* the edges and internal vertices of  $C[u, v]$ . For a chord  $e$ , let  $F(e)$  consist of  $e$  and all chords covering the end of  $e$ .

Select a chord  $e_1$  so that  $|F(e_1)| \geq |F(e)|$  for every chord  $e$ . For  $j > 1$ , let  $e_j$  be the first chord encountered moving forward from  $e_{j-1}$  that does not overlap  $e_{j-1}$  or  $e_1$ ; if no such chord exists, then stop and set  $\alpha = j - 1$ . Note that  $F(e_i) \cap \{e_1, \dots, e_\alpha\} = \{e_i\}$  for each  $i$  and that the sets  $F(e_1), \dots, F(e_\alpha)$  are pairwise disjoint. The selected edges  $\{e_1, \dots, e_\alpha\}$  form a *greedy chord system* for  $G$  (see Figure 1, which also includes notation used in Theorem 5). Given a greedy chord system beginning with  $e_1$ , let  $v_1$  be the start of  $e_1$ , and let the vertices of  $C$  be  $v_1, \dots, v_n$  in forward order.




Figure 1: A greedy chord system

From a greedy chord system, we will build a large family of cycles with distinct lengths by using short cycles, long cycles, and cycles of intermediate lengths. The intermediate-length cycles are formed from the long cycles by replacing portions of  $C$  with chords.

**Theorem 5** *Let  $G$  be a graph consisting of an  $n$ -cycle  $C$  plus  $q$  chords of length  $l$ , where  $l < n/2$ . The size  $s(G)$  of the cycle spectrum of  $G$  is at least  $(q-1)/2$  when  $l$  is even and at least  $(q-1-\frac{q}{l})/2$  when  $l$  is odd.*

**Proof.** Consider a greedy chord system  $e_1, \dots, e_\alpha$ . Let  $F' = F(e_1)$ . Let  $w$  be the end of the last chord in  $F'$  (see Figure 1). Let  $F^*$  be the set of chords not in  $\bigcup_{i=1}^\alpha F(e_i)$ ; since none of these chords overlaps  $e_\alpha$ , each overlaps  $e_1$ . If  $F^* \neq \emptyset$ , then let  $e^*$  be the first chord of  $F^*$  following  $e_\alpha$  (see Figure 1).

When  $\alpha = 1$ , we have  $|F'| + |F^*| = q$ . If also  $F^* = \emptyset$ , then  $|F'| = q$ . Otherwise,  $F^* \subseteq F(e^*) - \{e_1\}$ , so  $|F^*| \leq |F(e^*)| - 1 \leq |F'| - 1$ . Hence  $|F'| - 1 \geq (q-1)/2$ . Lemma 3 now yields  $v_1, w$ -paths of at least  $(q-1)/2$  lengths that combine with  $C[w, v_1]$  to form cycles of at least  $(q-1)/2$  lengths. Hence we may assume  $\alpha \geq 2$ .

For  $\alpha \geq 2$ , we begin by using  $F^*$  to obtain at least  $(|F^*| - 1)/2$  short cycle lengths. We may assume  $|F^*| \geq 2$ . Define  $j$  by  $e^* = v_j v_{j+l-n}$ . Through each chord  $v_k v_{k+l-n}$  in  $F^* - \{e^*\}$ , consider two cycles. One uses  $v_k v_{k+l-n}$  and  $e^*$  and the two paths  $C[v_j, v_k]$  and  $C[v_{j+l-n}, v_{k+l-n}]$  that each have length  $k - j$  (see Figure 1). The other uses  $v_k v_{k+l-n}$  and  $e_1$  and the two paths  $C[v_k, v_1]$  and  $C[v_{k+l-n}, v_{1+l}]$  that each have length  $n - k + 1$ . The lengths of these cycles are  $2(k - j + 1)$  and  $2(n - k + 2)$ ; their minimum is at most  $n - j + 3$ .

Taking the shorter for each  $k$ , we obtain  $|F^*| - 1$  cycles having length at most  $n - j + 3$ , with each length occurring at most twice. This yields a set  $Q$  of  $(|F^*| - 1)/2$  values bounded by  $n - j + 3$ . Since  $v_j$  is between the end of  $e_\alpha$  and  $v_n$ , we have  $j \geq 1 + \alpha l$ , and values in  $Q$  are bounded by  $n - \alpha l + 2$ . Since  $\alpha \geq 2$ , these values are at most  $n - \alpha(l - 1)$ .

With  $\alpha \geq 2$ , let  $z$  be the end of  $e_2$  (see Figure 1), and say that a cycle in  $G$  is *long* if it contains  $C[z, v_1]$  and has length at least  $n - 2(l - 1) + 1$ . Let  $R$  be the set of lengths of long cycles, and let  $\rho = |R|$ .

From the long cycles in  $G$ , we construct shorter cycles. Since long cycles contain  $C[z, v_1]$ , they contain all edges of  $C$  covered by any of  $e_3, \dots, e_\alpha$ . These chords are pairwise non-overlapping and can replace parts of long cycles. Each such replacement yields  $\rho$  distinct lengths (within an interval of  $2(l - 1)$  values), shorter by  $l - 1$  than the previous set of lengths.

The set  $R$  and the  $\alpha - 2$  sets of size  $\rho$  produced by using  $e_3, \dots, e_\alpha$  successively to reduce lengths together form  $\alpha - 1$  sets of size  $\rho$ . Since each set lies in an interval of length  $2(l - 1)$ , each value appears in at most two of the sets. Also, the top part of  $R$  (values exceeding  $n - (l - 1)$ ) and the bottom part of the last translation (values at most  $n - (\alpha - 1)(l - 1)$ ) appear only once. Let  $R'$  be the union of those two sets. Since every value in  $R$  is above  $n - (l - 1)$  or at most  $n - (l - 1)$ , we have  $|R'| = \rho$ . Including also  $R'$ , we now have  $\alpha$  sets of size  $\rho$ , with each value appearing in at most two of them.

Hence the union contains at least  $\alpha\rho/2$  cycle lengths, all at least  $n - \alpha(l - 1) + 1$  (which exceeds  $\max Q$ ). Thus  $s(G) \geq (\alpha\rho + |F^*| - 1)/2$ . It remains to study this quantity.

The greedy choice of  $e_1$  yields  $|F'| \geq \left\lceil \frac{q-|F^*|}{\alpha} \right\rceil$ . To obtain a lower bound on  $\alpha\rho$ , we compare  $\rho$  to  $|F'|$ . Let  $G'$  be the induced subgraph of  $G$  consisting of  $C[v_1, w]$  and the chords in  $F'$ . Since these chords are pairwise overlapping, Lemma 3 yields  $v_1, w$ -paths in  $G'$  with  $|F'| - 1$  distinct lengths. Furthermore, there are at least  $|F'|$  distinct lengths unless  $l$  is odd,  $|F'| \geq (l+3)/2$ , and the starts of the chords in  $F'$  are consecutive along  $C$ .

If  $|F'| = 1$ , then the greedy choice of  $e_1$  implies that the chords are pairwise noncrossing and  $s(G) = q + 1$ . We may thus assume  $|F'| > 1$  and  $w \neq v_{l+1}$ , so every  $v_1, w$ -path in  $G'$  has length at least 2. Adding  $C[w, v_1]$  to  $v_1, w$ -paths of distinct lengths in  $G'$  creates cycles of distinct lengths in  $G$ . Since each such cycle contains  $C[w, v_1]$ , which has at least  $n - 2l + 1$  edges, these cycles are long.

Thus when  $l$  is even, we have shown that  $\rho \geq |F'|$ . In this case

$$s(G) \geq \frac{\alpha\rho}{2} + \frac{|F^*| - 1}{2} \geq \frac{q - |F^*|}{2} + \frac{|F^*| - 1}{2} = \frac{q - 1}{2}.$$

If  $l$  is odd, then  $\frac{\alpha\rho}{2} \geq \left\lceil \frac{q-|F^*|}{\alpha} \right\rceil$  still holds if  $|F'| > \left\lceil \frac{q-|F^*|}{\alpha} \right\rceil$ , since  $\rho \geq |F'| - 1$ . Hence we may assume  $|F'| \geq \left\lceil \frac{q-|F^*|}{\alpha} \right\rceil$ . If  $\rho \geq |F'|$  fails, then Lemma 3 implies that  $|F'| \geq (l+3)/2$  and that the chords in  $F'$  are consecutive. Now  $R$  consists of the  $|F'| - 1$  values from  $n$  through  $n - 2|F'| + 4$  whose difference from  $n$  is even. We consider two cases, depending on whether  $e_2$  overlaps some chord in  $F'$ .

**Case 1:**  $e_2$  overlaps no chord in  $F'$ . Here  $e_2$ , like  $e_3, \dots, e_\alpha$ , can be used to reduce cycle lengths by  $l - 1$ . Since  $|F'| \geq (l+3)/2$ , the long cycle lengths include  $n, n - 2, \dots, n - (l - 1)$ ; there are  $(l + 1)/2$  of them. After using each of  $e_2, \dots, e_\alpha$  to reduce the lengths by  $l - 1$ , we obtain all values with the same parity as  $n$  down to  $n - \alpha(l - 1)$ . The smallest may equal  $\max Q$ . We keep  $\frac{1}{2}\alpha(l - 1)$  cycle lengths, each at least  $n - \alpha(l - 1) + 2$ .

If  $\alpha \geq q/l$ , then  $\frac{1}{2}\alpha(l - 1) \geq \frac{1}{2}q(1 - \frac{1}{l}) \geq \frac{1}{2}(q - |F^*| - \frac{q}{l})$ . If  $\alpha < q/l$ , then we use  $l \geq |F'| = \left\lceil \frac{q-|F^*|}{\alpha} \right\rceil$  to compute

$$\frac{1}{2}\alpha(l - 1) \geq \frac{1}{2}(|F'| - 1)\alpha \geq \frac{1}{2}(q - |F^*| - \alpha) > \frac{1}{2}\left(q - |F^*| - \frac{q}{l}\right).$$

Adding the  $(|F^*| - 1)/2$  short lengths yields at least the desired number of lengths.

**Case 2:**  $e_2$  overlaps some chord in  $F'$ . Since the chords in  $F'$  are consecutive, this case requires that  $e_2$  starts just before the end of some chord  $e'$  in  $F'$ . Let  $v'$  be the start of  $e'$ . The cycle consisting of  $e_2$  and  $e'$ , the edge they both cover, and the path  $C[z, v']$  (see Figure 1) has length  $n - 2(l - 1) + 2$ ; hence it is a long cycle. We obtain  $\rho \geq |F'|$  unless this length already appears among those generated from Lemma 3, which requires  $2|F'| - 4 \geq 2(l - 1) - 2$ , so  $|F'| \geq l$ . Since  $|F'| \leq l$ , equality holds.

As noted above, already  $n, n-2, \dots, n-2(l-2) \in R$ . Lowering the bottom half of them by  $l-1$  exactly  $\alpha-2$  times yields  $\frac{1}{2}\alpha(l-1)$  distinct cycle lengths. The least of them is  $n-\alpha(l-1)+2$ . This is exactly the situation we obtained in Case 1, so the same computation completes the proof.  $\square$

**Theorem 6** *If  $G$  is an  $n$ -vertex Hamiltonian graph with  $m$  edges, then  $s(G) > \sqrt{p}-\frac{1}{2}\ln p-1$ , where  $p = m - n$ .*

**Proof.** Let  $C$  be a spanning cycle in  $G$ . Let  $L$  be the set of lengths of chords of  $C$  in  $G$ , and let  $t = |L|$ . For each  $l \in L$ , we obtain two lengths of cycles in  $G$ ; they are  $l+1$  and  $n-l+1$  if  $l < n/2$  (using one chord of length  $l$ ), and they are  $n/2+1$  and  $n$  if  $l = n/2$ . Hence  $s(G) \geq 2t$ , which suffices if  $t \geq \frac{1}{2}\sqrt{p}$ . We may therefore assume that  $2t < \sqrt{p}$ .

For  $l \in L$ , let  $q_l$  be the number of chords of length  $l$ . By Theorem 5, when  $l < n/2$  there are at least  $\frac{l-1}{2l}q_l - \frac{1}{2}$  lengths of cycles using only edges of  $C$  and chords of length  $l$ . The lower bound also holds when  $l = n/2$ , since then the chords are pairwise overlapping and Lemma 3 applies, and always  $q_l - 1 > \frac{l-1}{2l}q_l - \frac{1}{2}$ .

We may assume that  $\frac{l-1}{2l}q_l - \frac{1}{2} \leq \sqrt{p} - \frac{1}{2}\ln p - 1$  for odd  $l \in L$ , and  $\frac{1}{2}q_l - \frac{1}{2} \leq \sqrt{p} - \frac{1}{2}\ln p - 1$  for even  $l \in L$ . Thus  $q_l \leq (\sqrt{p} - \frac{1}{2}\ln p - \frac{1}{2})c_l$ , where  $c_l = 2$  when  $l$  is even and  $c_l = 2 + \frac{2}{l-1}$  when  $l$  is odd. We obtain a contradiction by showing that these bounds on  $q_l$  sum to less than  $p$ . In light of the form of  $c_l$ , it suffices to prove this when all values in  $L$  are odd. The bound is now the worst when  $L$  consists of the first  $t$  positive odd numbers. We compute

$$\begin{aligned} p = \sum_{l \in L} q_l &\leq \sum_{l \in L} \left( \sqrt{p} - \frac{1}{2}\ln p - \frac{1}{2} \right) \left( 2 + \frac{2}{l-1} \right) \leq \left( \sqrt{p} - \frac{1}{2}\ln p - \frac{1}{2} \right) \left[ 2t + \sum_{i=1}^t \frac{1}{i} \right] \\ &< \left( \sqrt{p} - \frac{1}{2}\ln p - \frac{1}{2} \right) [\sqrt{p} + (1 + \ln t)] < \left( \sqrt{p} - \frac{1}{2}\ln p - \frac{1}{2} \right) [\sqrt{p} + \frac{1}{2}\ln p + (1 - \ln 2)] \\ &= p - \frac{1}{4}(\ln p)^2 - (\ln 2 - \frac{1}{2})\sqrt{p} - \frac{1}{4}(3 - \ln 4)\ln p - \frac{1}{2}(1 - \ln 2) < p. \end{aligned}$$

The contradiction completes the proof.  $\square$

## References

- [1] D. Bauer and E. F. Schmeichel, Hamiltonian degree conditions which imply a graph is pancyclic. *J. Combin. Theory (B)* 48 (1990), 111-116.
- [2] B. Bollobás and A. Thomason, Weakly pancyclic graphs. *J. Comb. Th. (B)* 77 (1999), 121-137.
- [3] J.A. Bondy, Pancyclic Graphs. *J. Combin. Theory (B)* 11 (1971), 80-84.

- [4] J.A. Bondy, Longest paths and cycles in graphs of high degree. Res. Rep. No. CORR 80-16, Univ. Waterloo, Waterloo, ON, 1980.
- [5] V. Chvátal, On Hamilton's ideals. *J. Combin. Theory (B)* 12 (1972), 163-168.
- [6] R.C. Entringer and E.F. Schmeichel, Edge conditions and cycle structure in bipartite graphs. *Ars Combinatoria* 26 (1988), 229-232.
- [7] P. Erdős, Some of my favourite problems in various branches of combinatorics. *Matematiche (Catania)* 47 (1992), 231-240.
- [8] P. Erdős, R. Faudree, C. Rousseau, and R. Schelp, The number of cycle lengths in graphs of given minimum degree and girth. *Discrete Math.* 200 (1999), 55-60.
- [9] G. Fan, New sufficient conditions for cycles in graphs. *J. Combin. Th. (B)* 37 (1984), 221-227.
- [10] M. Ferrara, A. Harris, and M. Jacobson, Cycle lengths in Hamiltonian graphs with a pair of vertices having large degree sum. *Graphs and Combinatorics* 26 (2010), 215-223.
- [11] R. Gould, P. Haxell, and A. Scott, A note on cycle lengths in graphs. *Graphs and Combinatorics* 18 (2002), 491-498.
- [12] A. Pnueli, A. Lempel, and S. Even, Transitive orientation of graphs and identification of permutation graphs. *Canad. J. Math.* 23 (1971), 160-175.
- [13] E.F. Schmeichel and S.L. Hakimi, A cycle structure theorem for Hamiltonian graphs. *J. Combin. Theory (B)* 45 (1988), 99-107.
- [14] E.F. Schmeichel and J. Mitchem, Bipartite graphs with cycles of all even lengths. *J. Graph Theory* 6 (1982), 429-439.
- [15] B. Sudakov and J. Verstraëte, Cycle lengths in sparse graphs, *Combinator.* 28 (2008), 357-372.
- [16] O. Ore, Note on Hamilton circuits. *Amer. Math. Monthly* 67 (1960), 55.